Click to verify



```
a. The balanced chemical equation for the complete combustion of Methanol is:-2CH3OH(l)+3O2(g) \rightarrow 8CO2(g)+4H2O(l)MethanolairCarbonWater(Oxygen)dioxide 01 We need
to find out the balanced chemical equation for the combustion of methanol. 02 Alcohols are combustion of methanol. 02 Alcohols are combustion of methanol of motorial equation for the presence of a hydrocarbon chain. Alcohols totally combust in the presence of oxygen to produce carbon monoxide or
carbon (ash). Combustion reaction of Methanol: -2CH3OH(l)+3O2(g) -2CO2(g)+4H2O(l)MethanolairCarbonWater(Oxygen)dioxide 03 We need to find out the balanced chemical equation for the complete combustion of 2-butanol. 04 Alcohols are combustion of 2-butanol.
to produce carbon dioxide and water. They did not combust completely when oxygen is scant, delivering water and either carbon monoxide or carbon (ash). Combustion reaction of 2-butanol: -2CH3-CH-OHCH2-CH3(l)+12O2(g) \rightarrow8CO2(g)+10H2O(l)2-butanolairCarbonWater(Oxygen)dioxide Unlock Step-by-Step Solutions & Ace Your Exams! Full
Textbook Solutions Get detailed explanations and key concepts Unlimited Al creation Al flashcards, exams and more... Ads-free access To over 500 million students worldwide already upgrade their learning with Vaia! How can financial brands set
 themselves apart through visual storytelling? Our experts explain how.Learn MoreThe Motorsport Images Collections captures events from 1895 to today's most recent coverage. Discover The Collection Curated, compelling, and worth your time. Explore our latest gallery of Editors' Picks. Browse Editors' Favorites How can financial brands set
themselves apart through visual storytelling? Our experts explain how.Learn MoreThe Motorsport Images Collections captures events from 1895 to today's most recent coverage. Discover The Collection today's most recent coverage. Discover The Collection today's most recent coverage.
themselves apart through visual storytelling? Our experts explain how.Learn MoreThe Motorsport Images Collections captures events from 1895 to today's most recent coverage. Discover The Collection Curated, compelling, and worth your time. Explore our latest gallery of Editors' Picks. Browse Editors' Favorites Ch.18 - Free Energy and
ThermodynamicsPrevious problemNext problemNext problemNext fCO} 2 + \text{O} 2 \rightarrow \text{O} 2 + \text{H} 2\text{O} 2 + \text{H} 2\text{O} 2 + \text{H} 2\text{O} 2 + \text{D} 3\text{O} 3 + \text{D} 3\text{O} 3 + \text{D} 3\text{O} 3 + \text{D} 3\text{D} 3\tex
2\text{CH}_3\text{OH} + 3\text{O}_2 \rightarrow 2\text{CO}_2 + 4\text{H}_2\text{C}_1\text{rxn}} \) using the formula: \(\Delta H^\circ_{\text{rxn}} - \sum \Delta H^\circ_{\text{f,reactants}} \). Calculate \(\Delta H^\circ_{\text{rxn}} - \sum \Delta H^\circ_{\text{f,reactants}} \).
S^\circ \{ x^* \} \) using standard molar entropies: \\ \Delta G^\circ \{ x^* \} = S^\circ \{ x^* \} \) using standard molar entropies: \\ \Delta G^\circ \{ x^* \} = S^\circ \{ x^* \} \) using standard molar entropies: \\ \Delta G^\circ \{ x^* \} = S^\circ \{ x^* \} \) using standard molar entropies: \\ \Delta G^\circ \{ x^* \} = S^\circ \{ x^* \} \) using standard molar entropies: \\ \Delta G^\circ \{ x^* \} = S^\circ \{ x^* \} \) using standard molar entropies: \\ \Delta G^\circ \{ x^* \} = S^\circ \{ x^* \} \) using standard molar entropies: \\ \Delta G^\circ \{ x^* \} = S^\circ \{ x^* \} \) using standard molar entropies: \\ \Delta G^\circ \{ x^* \} = S^\circ \{ x^* \} \) using standard molar entropies: \\ \Delta G^\circ \{ x^* \} = S^\circ \{ x^* \} \) using standard molar entropies: \\ \Delta G^\circ \{ x^* \} = S^\circ \{ x^* \} \) using standard molar entropies: \\ \Delta G^\circ \{ x^* \} = S^\circ \{ x^* \} \) using standard molar entropies: \\ \Delta G^\circ \{ x^* \} = S^\circ \{ x^* \} \) using standard molar entropies: \\ \Delta G^\circ \{ x^* \} = S^\circ \{ x^* \} \) using standard molar entropies: \\ \Delta G^\circ \{ x^* \} = S^\circ \{ x^* \} \) using standard molar entropies: \\ \Delta G^\circ \{ x^* \} = S^\circ \{ x^* \} \) using standard molar entropies: \\ \Delta G^\circ \{ x^* \} = S^\circ \{ x^* \} \) using standard molar entropies: \\ \Delta G^\circ \{ x^* \} = S^\circ \{ x^* \} \) using standard molar entropies: \\ \Delta G^\circ \{ x^* \} = S^\circ \{ x^* \} \) using standard molar entropies: \\ \Delta G^\circ \{ x^* \} = S^\circ \{ x^* \} \) using standard molar entropies: \\ \Delta G^\circ \{ x^* \} = S^\circ \{ x^* \} \) using standard molar entropies: \\ \Delta G^\circ \{ x^* \} = S^\circ \{ x^* \} \) using standard molar entropies: \\ \Delta G^\circ \{ x^* \} = S^\circ \{ x^* \} \) using standard molar entropies: \\ \Delta G^\circ \{ x^* \} = S^\circ \{ x^* \} \
spontaneity by checking if \(\Delta G^\circ \\text{rxn}\) < 0 \). This video solution was recommended by our tutors as helpful for the problem above. OHere are the essential concepts you must grasp in order to answer the question correctly. Combustion reactions are exothermic processes where a substance reacts with oxygen to produce heat, light,
carbon dioxide, and water. In the case of methanol (CH3OH), the balanced equation for its combustion involves the complete reaction with oxygen to yield carbon dioxide (CO2) and water (H2O). Understanding how to balance these equations is crucial for stoichiometric calculations and thermodynamic assessments. 02:24Thermodynamics involves
the study of energy changes in chemical reactions. ΔH° (enthalpy change) indicates the heat absorbed or released, ΔS° (entropy change) measures the disorder or randomness in the system, and ΔG° (Gibbs free energy change) measures the disorder or randomness in the system, and ΔG° (Gibbs free energy change) measures the disorder or randomness in the system, and ΔG° (Gibbs free energy change) measures the disorder or randomness in the system, and ΔG° (Gibbs free energy change) measures the disorder or randomness in the system, and ΔG° (Gibbs free energy change) measures the disorder or randomness in the system, and ΔG° (Gibbs free energy change) measures the disorder or randomness in the system, and ΔG° (Gibbs free energy change) measures the disorder or randomness in the system, and ΔG° (Gibbs free energy change) measures the disorder or randomness in the system, and ΔG° (Gibbs free energy change) measures the disorder or randomness in the system, and ΔG° (Gibbs free energy change) measures the disorder or randomness in the system of the
conditions, which is essential for evaluating the combustion of methanol. 01:18First Law of ThermodynamicsThe spontaneity of a reaction refers to its ability to occur without external intervention. It is determined by the sign of AG°; if AG° is negative, the reaction is spontaneous. Factors influencing spontaneity include enthalpy and entropy changes,
where a highly exothermic reaction with a significant increase in entropy is likely to be spontaneous, as is the case with the combustion of methanol. 04:20Previous problemNext problemUse data from Appendix IIB to calculate \Delta S^{\circ}rxn for each of the reactions. In each case, try to rationalize the sign of \Delta S^{\circ}rxn. d. N2O4(g) + 4 H2(g) \rightarrow N2(g) + 4 H2(g) + 4 H2(g) \rightarrow N2(g) + 4 H2(g) \rightarrow N2(g) + 4 H2(g) \rightarrow N2(g) + 4 H2(g) 
H2O(g)Find ΔS° for the formation of CH2Cl2(g) from its gaseous elements in their standard states. Rationalize the sign of ΔS°. What is ΔS° for the reaction between nitrogen gas and fluorine gas to form nitrogen trifluoride gas, and how can the sign of ΔS°. What is ΔS° for the reaction between nitrogen gas and fluorine gas to form nitrogen trifluoride gas, and how can the sign of ΔS°. What is ΔS° for the reaction between nitrogen gas and fluorine gas to form nitrogen gas and fluorine gas.
dioxide and water. Write a balanced equation for photosynthesis and calculate ΔH°rxn, ΔS°rxn, and ΔG°rxn at 25 °C. Is photosynthesis spontaneous. If the reaction is not spontaneous, would a change in temperature make it spontaneous?
If so, should the temperature be raised or lowered from 25 °C? b. NH4Cl(s) \rightarrow HCl(g) + Fe2O3(s) \rightarrow 2 Fe(s) + 3 H2O(g)For each reaction is spontaneous. If the reaction is not spontaneous, would a change in temperature make it spontaneous? If
so, should the temperature be raised or lowered from 25 °C? a. N2O4(g) \rightarrow 2 NO2(g) In order to continue enjoying our site, we ask that you confirm your identity as a human. Thank you very much for your cooperation. InChI=1S/CH4O/c1-2/h2H,1H3InChIKey=OKKJLVBELUTLKV-UHFFFAOYSA-NSMILESCOCanonical SMILESCOCANONICAL SMI
Names for this SubstanceMethanolCarbinolMethyl alcoholMethyl alcoholMethyl hydroxideMonohydroxymethaneDeleted or Replaced CAS Registry Numbers54841-71-3, 1173023-83-0, 1196157-60-4 CH3OH; simplest possible alcohol Not to be confused with menthol, methanol, methanol, methanol, methanol, methanol, methanol, methanol, methanol, methanol Space filling model of methanol
Ball and stick model of methanol A sample of methanol Names Preferred IUPAC name Methanol[1] Other names CarbinolColumbian spirits HydroxymethaneMeOHMethyl hydroxideMethylic alcoholMethylolMethylene hydrate, primary alcoholPyroligneous spiritWood alcoholWood naphthaWood spirit Identifiers CAS
Number 67-56-1 Y 3D model (JSmol) Interactive image Beilstein Reference 1098229 ChEBI CHEBI:17790 Y ChEMBL 14688 Y ChemSpider 864 Y ECHA InfoCard 100.000.599 EC Number 200-659-6 Gmelin Reference 449 KEGG D02309 Y MeSH Methanol PubChem CID 887 RTECS number PC1400000 UNII Y4S76JWI15 Y UN number 1230
CompTox Dashboard (EPA) DTXSID2021731 InChI InChI=1S/CH4O/c1-2/h2H,1H3 YKey: OKKJLVBELUTLKV-UHFFFAOYAX SMILES CO Properties Chemical formula CH3OH Molar mass 32.042 g·mol-1 Appearance Colourless liquid Odor Faint and similar to ethanol Density
0.792 g/cm3[2] Melting point -97.6 °C (-143.7 °F; 175.6 K) Boiling point 64.7 °C (148.5 °F; 337.8 K) Solubility in water miscible log P -0.69 Vapor pressure 13.02 kPa (at 20 °C) Acidity (pKa) 15.5[3] Conjugate acid Methyloxonium[4] Conjugate base Methanolate[5] Magnetic susceptibility (\chi) -21.40\cdot10-6 cm3/mol Refractive index (nD) 1.33141[6]
 Viscosity 0.545 mPa·s (at 25 °C)[7] Dipole moment 1.69 D Thermochemistry Heat of combustion, higher value (HHV) 725.7 kJ/mol, 173.4 kcal/mol, 5.77 kcal/g Hazards[12][13] Occupational safety and health (OHS/OSH): Main hazards Methanol and its vapours are flammable. Moderately toxic for small animals - Highly toxic to large animals and
humans (in high concentrations) - May be fatal/lethal or cause blindness and damage to the liver, kidneys, and heart if swallowed - Toxicity effects from repeated over exposure have an accumulative effect on the central nervous system, especially the optic nervous system.
days after exposure[9] GHS labelling: Pictograms [8] Signal word Danger[8] Hazard statements H225, H301, H302, H305, H311, H331, H370[8] Precautionary statements P210, P233, P240, P241, P242, P243, P260, P264, P270, P271, P280, P301+P330+P331, P302+P352, P303+P361+P353, P304+P340, P305+P351+P338, P307+P311, P310, P31
P311, P312, P337+P313, P361, P363, P361, P363, P361, P363, P364 to 285 K) Autoignition temperature 470 °C (878 °F; 743 K)[15] 385 °C (725 °F; 658 K)[16] Explosive limits 6-36%[10] Lethal dose or concentration (LD, LC): LD50 (median dose) 5628 mg/kg
(rat, oral)7300 mg/kg (mouse, oral)12800 mg/kg (rat, oral)14200 mg/kg (rat, oral)14200 mg/kg (rat, oral)14200 mg/kg (rat, oral)1280 mg/kg (rat, oral)14200 mg/kg
mg/m3) ST 250 ppm (325 mg/m3) [skin][10] IDLH (Immediate danger) 6000 ppm[10] Safety data sheet (SDS) [1] Related compounds Related compounds Related compounds Related compounds MethanethiolSilanolEthanol (data page) Except where otherwise noted, data are given for materials in their standard state (at 25 °C [77 °F], 100 kPa).
N verify (what is YN?) Infobox references Chemical compound Methanol (also called methyl alcohol, with the chemical formula CH3OH (a methyl group linked to a hydroxyl group, often abbreviated as MeOH). It is a light, volatile, colorless and
flammable liquid with a distinctive alcoholic odor similar to that of ethanol (potable alcohol), but is more acutely toxic than the latter.[17] Methanol acquired the name wood alcohol because it was once produced through destructive distillation of wood. Today, methanol is mainly produced industrially by hydrogenation of carbon monoxide.[18]
Methanol consists of a methyl group linked to a polar hydroxyl group. With more than 20 million tons produced annually, it is used as a precursor to other commodity chemicals, including formaldehyde, acetic acid, methyl tert-butyl ether, methyl benzoate, anisole, peroxyacids, as well as a host of more specialized chemicals.[18] Small amounts of
methanol are present in normal, healthy human individuals. One study found a mean of 4.5 ppm in the exhaled breath of test subjects [19] The mean endogenous methanol in humans of 0.45 g/d may be metabolized from pectin found in fruit; one kilogram of apple produces up to 1.4 g of pectin (0.6 g of methanol.)[20] Methanol is produced by
anaerobic bacteria and phytoplankton.[21][22] Methanol is also found in abundant quantities in star-forming regions of space and is used in astronomy as a marker for such regions. It is detected through its spectral emission lines.[23] In 2006, astronomers using the MERLIN array of radio telescopes at Jodrell Bank Observatory discovered a large
cloud of methanol in space 0.463 terametres (288 million miles) across.[24][25] In 2016, astronomers detected methanol in a planet-forming disc around the young star TW Hydrae using the Atacama Large Millimeter Array radio telescope.[26] In their embalming process, the ancient Egyptians used a mixture of substances, including methanol, which
they obtained from the pyrolysis of wood. Pure methanol, however, was first isolated in 1661 by Robert Boyle, when he produced it via the distillation of buxus (boxwood).[27] It later became known as "pyroxylic spirit". In 1834, the French chemists Jean-Baptiste Dumas and Eugene Peligot determined its elemental composition.[28] They also
introduced the word "methylène" to organic chemistry, forming it from Greek methy = "alcoholic liquid" + hylē = "forest, wood, timber, material". "Methylène" designated a "radical" that was about 14% hydrogen by weight and contained one carbon atom. This would be CH2, but at the time carbon was thought to have an atomic weight only six times
that of hydrogen, so they gave the formula as CH.[28] They then called wood alcohol (l'esprit de bois) "bihydrate de méthylène" (bihydrate de méthylène" (bihydrate de méthylène", and was then applied to describe "methyl alcohol". This was
shortened to "methanol" in 1892 by the International Conference on Chemical Nomenclature, [29] The suffix -yl, which, in organic chemistry, forms names of carbon groups, is from the word methyl. French chemist Paul Sabatier presented that
carbon dioxide and hydrogen could be reacted to produce methanol.[9] German chemists Alwin Mittasch and Mathias Pier, working for Badische-Anilin & Soda-Fabrik (BASF), developed a means to convert synthesis gas (a mixture of carbon monoxide, carbon dioxide, and hydrogen) into methanol and received a patent. According to Bozzano and
Manenti, BASF's process was first utilized in Leuna, Germany in 1923. Operating conditions consisted of "high" temperatures (between 300 and 400 °C) and pressures (between 300 °C) and pressures (between 300 °C) and pressures (between 300 °C) and pressu
process used a chromium and manganese oxide catalyst with extremely vigorous conditions: pressures ranging from 50 to 220 atm, and temperatures up to 450 °C. Modern methanol production has been made more efficient through use of catalysts (commonly copper) capable of operating at lower pressures. The modern low pressure methanol (LPM)
process was developed by ICI in the late 1960s with the technology patent long since expired.[32] During World War II, methanol was used as a fuel in several German military rocket designs, under the name M-Stoff, and in a roughly 50/50 mixture with hydrazine, known as C-Stoff. The use of methanol as a motor fuel received attention during the oil
crises of the 1970s. By the mid-1990s, over 20,000 methanol flexible fuel vehicles (FFVs) capable of operating on methanol or gasoline were introduced in the US. In addition, low levels of methanol were blended in gasoline were introduced in the US. In addition, low levels of methanol were blended in gasoline were introduced in the US. In addition, low levels of methanol were blended in gasoline were introduced in the US. In addition, low levels of methanol were blended in gasoline were introduced in the US. In addition, low levels of methanol were blended in gasoline were introduced in the US. In addition, low levels of methanol were blended in gasoline were introduced in the US. In addition, low levels of methanol were blended in gasoline were introduced in the US. In addition, low levels of methanol were blended in gasoline were introduced in the US. In addition, low levels of methanol were blended in gasoline were introduced in the US. In addition, low levels of methanol were blended in gasoline were introduced in the US. In addition, low levels of methanol were blended in gasoline were introduced in the US. In addition, low levels of methanol were blended in gasoline were introduced in the US. In addition, low levels of methanol were blended in gasoline were introduced in the US. In addition, low levels of methanol were blended in gasoline were also were also
[34] Between the 1960s and 1980s methanol emerged as a precursor to the feedstock chemicals acetic anhydride process. Methanol is primarily converted to formaldehyde, which is widely used in many areas,
 especially polymers. The conversion entails oxidation: 2 CH3OH + O2 \rightarrow 2 CH2O + 2 H2O Acetic acid can be produced from methanol. The Cativa process converts methanol into acetic acid. [35] Methanol and isobutene are combined to give methyl tert-butyl ether (MTBE). MTBE is a major octane booster in gasoline. Condensation of methanol to
produce hydrocarbons and even aromatic systems is the basis of several technologies related to gas to liquids. These conversions are catalyzed by zeolites as heterogeneous catalysts. The MtG process was once
commercialized at Motunui in New Zealand.[36][37] The European Fuel Quality Directive allows fuel producers to blend up to 3% methanol, with an equal amount of cosolvent, with gasoline sold in European Fuel Quality Directive allows fuel producers to blend up to 3% methanol, with an equal amount of cosolvent, with gasoline sold in European Fuel Quality Directive allows fuel producers to blend up to 3% methanol, with an equal amount of cosolvent, with gasoline sold in European Fuel Quality Directive allows fuel producers to blend up to 3% methanol, with an equal amount of cosolvent, with gasoline sold in European Fuel Quality Directive allows fuel producers to blend up to 3% methanol, with an equal amount of cosolvent, with gasoline sold in European Fuel Quality Directive allows fuel producers to blend up to 3% methanol, with an equal amount of cosolvent, with gasoline sold in European Fuel Quality Directive allows fuel producers to blend up to 3% methanol, with an equal amount of cosolvent, with gasoline sold in European Fuel Quality Directive allows fuel producers to blend up to 3% methanol, with an equal amount of cosolvent, with gasoline sold in European Fuel Quality Directive allows fuel producers to blend up to 3% methanol and producers to 3% methanol and produc
Methanol is the precursor to most simple methylamines, methyl halides, and methyl esters are produced from methanol, including the transesterification of fats and production of biodiesel via transesterification of fats and produced from methanol, including the transesterification of fats and produced from methanol, including the transesterification of fats and produced from methanol, including the transesterification of fats and produced from methanol, including the transesterification of fats and produced from methanol, including the transesterification of fats and produced from methanol, including the transesterification of fats and produced from methanol, including the transesterification of fats and produced from methanol, including the transesterification of fats and produced from methanol, including the transesterification of fats and produced from methanol, including the transesterification of fats and produced from methanol fats and produced from methanol fats are produced from methanol fats and produced from methanol fats are pro
 Its energy density is, however, lower than methane, per kg. Its combustion energy density is 15.6 MJ/L. (LHV), whereas that of ethanol is 24 and gasoline is 33 MJ/L. Further advantages for methanol is 24 and gasoline is 33 MJ/L.
environments. The half-life for methanol in groundwater is just one to seven days, while many common gasoline components have half-lives in the hundreds of days (such as benzene at 10-730 days). Since methanol is miscible with water and biodegradable, it is unlikely to accumulate in groundwater is just one to seven days, while many common gasoline components have half-lives in the hundreds of days (such as benzene at 10-730 days).
Methanol fuel and methanol economy Methanol is occasionally used to fuel internal combustion engines. It burns forming carbon dioxide and water: 2 CH3OH + 3 O2 → 2 CO2 + 4 H2O Methanol fuel has been proposed for ground transportation. The chief advantage of a methanol economy is that it could be adapted to gasoline internal combustion
engines with minimum modification to the engines and to the infrastructure that delivers and stores liquid fuel. Its energy density, however, it is equivalent to super high-octane gasoline in horsepower, and most modern computer-controlled fuel injection systems can
already use it.[42] Methanol is an alternative fuel for ships that helps the shipping industry meet increasingly strict emissions of sulfur oxides (NOx), nitrogen oxides (NOx) and particulate matter. Methanol can be used with high efficiency in marine diesel engines after minor modifications using a small
amount of pilot fuel (dual fuel).[43][44] In China, methanol fuels industrial boilers, which are used extensively to generate heat and steam for various industrial applications and residential heating. Its use is displacing coal, which is under pressure from increasingly stringent environmental regulations.[45] Direct-methanol fuels are unique in their
low temperature, atmospheric pressure operation, which lets them be greatly miniaturized. [46] [47] This, combined with the relatively easy and safe storage and handling of methanol, may open the possibility of fuel cell-powered consumer electronics, such as laptop computers and mobile phones.
and boating stoves. Methanol burns well in an unpressurized burner, so alcohol stoves are often very simple, sometimes little more than a cup to hold fuel. This lack of complexity makes them a favorite of hikers who spend extended time in the wilderness. Similarly, the alcohol can be gelled to reduce risk of leaking or spilling, as with the brand
 "Sterno". Methanol is mixed with water and injected into high performance diesel and gasoline engines for an increase of power and a decrease in intake air temperature in a process known as denatured alcohol or methylated spirit. This was
commonly used during the US prohibition to discourage consumption of bootlegged liquor, and ended up causing several deaths.[49] It is sometimes used as a fuel in alcohol lamps, portable fire pits and camping stoves. Methanol is used as a fuel in alcohol lamps, portable fire pits and camping stoves.
coolant antifreeze in the early 1900s.[50] As of May 2018, methanol was banned in the EU for use in windscreen washing or defrosting due to its risk of human consumption[51][52] as a result of 2012 Czech Republic methanol poisonings.[53] In some wastewater treatment plants, a small amount of methanol is added to wastewater to provide a
 carbon food source for the denitrifying bacteria, which convert nitrates to nitrogen gas and reduce the nitrification of sensitive aquifers. Methanol is used as a destaining agent in polyacrylamide gel electrophoresis. Carbon monoxide and hydrogen react over a catalyst to produce methanol. Today, the most widely used catalyst is a mixture of copperation.
and zinc oxides, supported on alumina, as first used by ICI in 1966. At 5-10 MPa (50-100 atm) and 250 °C (482 °F), the reaction CO + 2 H2 → CH3OH is characterized by high selectivity (>99.8%). The production of synthesis gas from methane produces three moles of hydrogen for every mole of carbon monoxide, whereas the synthesis consumes only
two moles of hydrogen gas per mole of carbon monoxide. One way of dealing with the excess hydrogen is to inject carbon dioxide into the methanol synthesis reactor, where it, too, reacts to form methanol according to the equation CO2 + 3 H2 - CH3OH + H2O In terms of mechanism, the process occurs via initial conversion of CO into CO2, which is
then hydrogenated: [54] CO2 + 3 H2 \rightarrow CH3OH + H2O where the H2O byproduct is recycled via the water-gas shift reaction CO + 2 H2 \rightarrow CH3OH which is the same as listed above. In a process closely related to methanol production from synthesis gas, a feed of hydrogen and CO2 can be used
directly.[55] The main advantage of this process is that captured CO2 and hydrogen sourced from electrolysis could be used, removing the dependence on fossil fuels. The catalytic conversion of methane to methanol is effected by enzymes including methane monooxygenases. These enzymes are mixed-function oxygenases, i.e. oxygenation is coupled
with production of water[56] and NAD+:[57] CH4 + O2 + NADPH + H+ \rightarrow CH3OH + H2O + NAD+ Both Fe- and Cu-dependent enzymes have been undertaken to emulate this reactivity.[58][59] Methanol is more easily oxidized than is the feedstock methane, so the reactions tend not to
be selective. Some strategies exist to circumvent this problem. Examples include Shilov systems and Fe- and Cu-containing zeolites. [60] These systems do not necessarily mimic the mechanisms employed by metalloenzymes, but draw some inspiration from them. Active sites can vary substantially from those known in the enzymes. For example, a
dinuclear active site is proposed in the Fe-zeolite.[61] Global emissions of methanol by plants are estimated at between 180 and 250 million tons per year.[62] This is between two and three times larger than man-made industrial production of methanol. As of 2023, 0.2% of
 global methanol production is produced in ways that have relatively low greenhouse gas emissions; this is known as "green" methanol.[63] Most green methanol from biomass gasification and further converted into green methanol.[64] Another method of producing green methanol
 involves combining hydrogen, carbon dioxide, and a catalyst under high heat and pressure.[63] To be classified as green methanol, the hydrogen must be produced using renewable electricity,[63] and the carbon dioxide must be from carbon dioxide must be fr
specify that the carbon dioxide must be captured during the burning of bioenergy.[65] Methanol is available commercially in various purity grades A and AA. Both grade AA purity are 99.85% methanol by weight. Grade "AA" methanol contains trace amounts of
ethanol as well.[30] Methanol for chemical use normally corresponds to Grade AA. In addition to water, typical impurities include acetone and ethanol (which are very difficult to separate by distillation). UV-vis spectroscopy is a convenient method for detecting aromatic impurities. Water content can be determined by the Karl-Fischer titration.
Methanol is highly flammable. Its vapours are slightly heavier than air and can travel to a distant ignition source and ignite. Methanol flames are invisible in daylight. Main article: Methanol toxicity See also: List of methanol poisoning
incidents Ingesting as little as 10 mL (0.34 US fl oz) of pure methanol can cause permanent blindness by destruction of the optic nerve. 30 mL (1.0 US fl oz), i.e., 1-2 mL/kg body weight of pure methanol.[67] The reference dose for methanol is 0.5 mg/kg in a day.[68][69] Toxic
effects begin hours after ingestion, and antidotes can often prevent permanent damage. [66] Because of its similarities in both appearance and odor to ethanol is toxic by two mechanisms. First, methanol can be fatal due to effects on the
central nervous system, acting as a central nervous system depressant in the same manner as ethanol poisoning. Second, in a process initiated by the enzyme alcohol dehydrogenase in the liver.[71] Methanol is converted to formaldehyde
via alcohol dehydrogenase (ADH) and formaldehyde is converted to formate via ALDH proceeds completely, with no detectable formate via ALDH proceeds completely via ALDH proceeds comp
metabolic acidosis, among a variety of other metabolic disturbances. [73] Outbreaks of methanol poisoning have occurred primarily due to contaminated batches of alcohol contained a much higher concentration of methanol than
ethanol, which did not provide a sufficient dose of the antidote and caused it to be ineffective. This is more common in the developing world.[75] In 2013 more than 1700 cases nonetheless occurred in the United States. Those affected are often adult men.[76] Outcomes may be good with early treatment.[77] Toxicity to methanol was described as
early as 1856.[78] Because of its toxic properties, methanol is frequently used as a denaturant additive for ethanol manufactured for industrial uses. This addition of methanol exempts industrial ethanol (commonly known as "denaturant additive for ethanol manufactured for industrial uses. This addition of methanol manufactured for industrial uses. This addition of methanol exempts industrial uses.
(data page) Trimethyl carbinol ^ Nomenclature of Organic Chemistry: IUPAC Recommendations and Preferred Names 2013 (Blue Book). Cambridge: The Royal Society of Chemistry and Physics (86th ed.). Boca Raton,
 Florida: CRC Press. ISBN 0-8493-0486-5. Sallinger, P.; Long, F. A. (1960). "Acid Ionization Constants of Alcohols. II. Acidities of Some Substituted Methanols and Related Compounds". J. Am. Chem. Soc. 82 (4): 795-798. Bibcode:1960JAChS..82..795B. doi:10.1021/ja01489a008. Substituted Methanols and Related Compounds". J. Am. Chem. Soc. 82 (4): 795-798. Bibcode:1960JAChS..82..795B. doi:10.1021/ja01489a008.
original on 21 December 2018. Retrieved 21 December 2018. The conjugate base of a methanol. "Refractive Index.INFO - Refractive Index.INFO - R
database". refractive index.info. Archived from the original on 23 February 2017. Retrieved 14 February 2012. ^ González, Begoña (2007). "Density, dynamic viscosity, and derived properties of binary mixtures of methanol or ethanol with water, ethyl acetate, and methyl acetate at T = (293.15, 298.15, and 303.15) K". The Journal of Chemical
Thermodynamics. 39 (12): 1578-1588. Bibcode:2007JChTh...39.1578G. doi:10.1016/j.jct.2007.05.004. ^ a b c d "Methanol" (PDF). Lab Chem. Valtech. Archived 20 August 2018 at the Wayback Machine ^ a b c d NIOSH Pocket Guide to Chemical
 Hazards. "#0397". National Institute for Occupational Safety and Health (NIOSH). ^ a b c "Methanol". Immediately Dangerous to Life or Health Concentrations (IDLH). National Institute for Occupational Safety and Health (NIOSH). ^ a b c "Methanol". Immediately Dangerous to Life or Health Concentrations (IDLH). National Institute for Occupational Safety and Health (NIOSH). ^ a b c "Methanol". Immediately Dangerous to Life or Health Concentrations (IDLH). National Institute for Occupational Safety and Health (NIOSH). ^ a b c "Methanol". Immediately Dangerous to Life or Health Concentrations (IDLH). National Institute for Occupational Safety and Health (NIOSH). ^ a b c "Methanol". Immediately Dangerous to Life or Health Concentrations (IDLH). National Institute for Occupational Safety and Health Concentrations (IDLH). National Institute for Occupational Safety and Health Concentrations (IDLH). National Institute for Occupational Safety and Health Concentrations (IDLH). National Institute for Occupational Safety and Health Concentrations (IDLH). National Institute for Occupational Safety and Health Concentrations (IDLH). National Institute for Occupational Safety and Health Concentrations (IDLH). National Institute for Occupational Safety and Health Concentrations (IDLH). National Institute for Occupational Safety and Health Concentrations (IDLH). National Institute for Occupational Safety and Health Concentration (IDLH). National Institute for Occupational Safety and Health Concentration (IDLH). National Institute for Occupational Safety and Health Concentration (IDLH). National Institute for Occupational Safety and Health Concentration (IDLH). National Institute for Occupational Safety and Health Concentration (IDLH).
Control and Prevention. Archived from the original on 23 April 2009. Retrieved 3 August 2018. ^ "PubChem: Safety and Hazards - GHS Classification". National Library of Medicine. Archived from the original on 20 August 2018. ^ "Methanol Safe Handling Manual" (PDF)
Methanol Institute. 2017. p. 253. Archived (PDF) from the original on 20 December 2017. Retrieved 3 August 2018. ^ "Technical Information & Safe Handling Manual" (PDF). Methanol Institute. 2017. p. 243. Archived (PDF) from
the original on 20 December 2017. Retrieved 3 August 2018. A National Institute for Occupational Safety and Health Database: Methanol". Archived from the original on 23 April 2009. A a b c Fiedler, E.; Grossmann, G.; Burkhard Kersebohm, D.; Weiss, G.;
Witte, C. (2005). "Methanol". Ullmann's Encyclopedia of Industrial Chemistry. Weinheim: Wiley-VCH. doi:10.1002/14356007.a16 465. ISBN 978-3-527-30673-2. ^ Turner C (2006). "A longitudinal study of methanol in the exhaled breath of 30 healthy volunteers using selected ion flow tube mass spectrometry, SIFT-MS". Physiological Measurement. 27
(7): 637-48. Bibcode:2006PhyM...27..637T. doi:10.1088/0967-3334/27/7/007. PMID 16705261. S2CID 22365066. ^ Lindinger W (1997). "Endogenous production of fruit". Alcoholism: Clinical and Experimental Research. 21 (5): 939-43. doi:10.1111/j.1530-0277.1997.tb03862.x. PMID 9267548. ^ "Major Source of
 Methanol in the Ocean Identified". Woods Hole Oceanographic Institution. 10 March 2016. Archived from the original on 28 December 2018. Retrieved 30 March 2016. ^ Mincer, Tracy J.; Aicher, Athena C. (2016). "Methanol Production by a Broad Phylogenetic Array of Marine Phytoplankton". PLOS ONE. 11 (3): e0150820.
Bibcode:2016PLoSO..1150820M. doi:10.1371/journal.pone.0150820. PMC 4786210. PMID 26963515. ^ Brooks Hays (17 April 2015). "Why astronomers hate the lawn-mowing Roomba". Space Daily. Archived from the original on 27 April 2015. ^ "Upgraded MERLIN spies cloud of alcohol spanning 288 billion miles" (Press
release). Jodrell Bank Centre for Astrophysics. 19 April 2006. Archived from the original on 6 October 2014. Retrieved 5 September 2010. ^ "First Detection of Methyl Alcohol in a Planet-forming Disc". Archived from the original
on 22 June 2016. Retrieved 22 June 2016. ^ Boyle discusses the distillation of liquids from the wood of the box shrub in: Robert Boyle, The Sceptical Chymist (London, England: J. Cadwell, 1661), pp. 192-195. ^ a b A report on methanol to the French Academy of Sciences by J. Dumas and E. Péligot began during the Academy's meeting of 27 October
1834 and finished during the meeting of 3 November 1834. See: Procès-verbaux des séances de l'Académie, 10: 600-601. Available on: Gallica Archived 25 April 2015 at the Wayback Machine. The complete report appears in: J. Dumas and E. Péligot (1835) "Mémoire sur l'espirit de bois et sur les divers composés ethérés qui en proviennent" (Memoire sur l'espirit de bois et sur les divers composés ethérés qui en proviennent" (Memoire sur l'espirit de bois et sur les divers composés ethérés qui en proviennent (Memoire sur l'espirit de bois et sur les divers composés ethérés qui en proviennent (Memoire sur l'espirit de bois et sur les divers composés ethérés qui en proviennent (Memoire sur l'espirit de bois et sur les divers composés ethérés qui en proviennent (Memoire sur l'espirit de bois et sur les divers composés ethérés qui en proviennent (Memoire sur l'espirit de bois et sur les divers composés ethérés qui en proviennent (Memoire sur l'espirit de bois et sur les divers composés ethérés qui en proviennent (Memoire sur l'espirit de bois et sur les divers composés ethérés qui en proviennent (Memoire sur l'espirit de bois et sur les divers composés ethérés qui en proviennent (Memoire sur l'espirit de bois et sur les divers composés ethérés qui en proviennent (Memoire sur l'espirit de bois et sur les divers de l'espirit de l'espiri
on spirit of wood and on the various ethereal compounds that derive therefrom), Annales de chimie et de physique, 58: 5-74; from page 9: Nous donnerons le nom de méthylène (1) à un radical ... (1) Mεθυ, vin, et υλη, bois; c'est-à-dire vin ou liqueur spiritueuse du bois. (We will give the name methylene (1) to a radical ... (1) methy, wine, and hulē,
wood; that is, wine or spirit of wood.) ^ For a report on the International Conference on Chemical Nomenclature that was held in April 1892 in Geneva, Switzerland, see: Armstrong, Henry E (1892). "The International Conference on Chemical Nomenclature". Nature. 46 (1177): 56-9. Bibcode:1892Natur..46...56A. doi:10.1038/046056c0. Armstrong's
report is reprinted with the resolutions in English in: Armstrong, Henry (1892). "The International Conference on Chemical Nomenclature". The Journal of Analytical and Applied Chemistry. 6 (1177): 390-400. Bibcode: 1892Natur..46...56A. doi:10.1038/046056c0. p. 398: 15. The alcohols and the phenols are named after the hydrocarbon from which
they derive, terminated with the suffix of (ex. pentanol, pentenol, etc.). ^ a b Bozzano, Giulia; Manenti, Flavio (1 September 2016). "Efficient methanol synthesis: Perspectives, technologies and optimization strategies". Progress in Energy and Combustion Science. 56: 71-105. Bibcode: 2016PECS...56...71B. doi:10.1016/j.pecs.2016.06.001. ISSN 0360-
1285. Archived from the original on 25 September 2021. Retrieved 25 September 2021. ^ US 3326956 ^ Halderman, James D.; Martin, Tony (2009). Hybrid and alternative fuel vehicles. Pearson/Prentice Hall. ISBN 978-0-13-504414-8. ^ Ronald Smith (1 December 2011). "Methanol to Gasoline: A Private Report by the Process
 Economics Program" (PDF). Archived (PDF) from the original on 4 December 2019. Achieved 4 December 2019. Sunley, G. J.; Watson, D. J. (2000). "High productivity methanol carbonylation catalysis using iridium - The Cativa process for the manufacture of acetic acid". Catalysis Today. 58 (4): 293-307. doi:10.1016/S0920-5861(00)00263-7.
 Olsbye, U.; Svelle, S.; Bjorgen, M.; Beato, P.; Janssens, T. V. W.; Joensen, F.; Bordiga, S.; Lillerud, K. P. (2012). "Conversion of Methanol to Hydrocarbons: How Zeolite Cavity and Pore Size Controls Product Selectivity". Angew. Chem. Int. Ed. 51 (24): 5810-5831. doi:10.1002/anie.201103657. hdl:2318/122770. PMID 22511469. S2CID 26585752.
Tian, P.; Wei, Y.; Ye, M.; Liu, Z. (2015). "Methanol to Olefins (MTO): From Fundamentals to Commercialization". ACS Catal. 5 (3): 1922-1938. doi:10.1021/acscatal.5b00007. "Biodiesel - METHANOL INSTITUTE". METHANOL INSTITUTE. April 2024. Retrieved 4 June 2024. "Biodiesel - METHANOL INSTITUTE". METHANOL INSTITUTE".
INSTITUTE. Archived from the original on 25 March 2018. Archived from the original on 25 March 2018. Archived 16 May 2016 at the Portuguese Web Archived 17 March 2018. Archived from the original on 25 March 2018.
Malcolm Pirnie, Inc., January 1999. ^ "Methanol Wins". National Review. December 2011. Archived from the original on 7 June 2022. Retrieved 7 June 2022. Retrieved 7 June 2021. Archived from the original on 7 June 2022. National Review. December 2011. Archived from the original on 7 June 2022. Retrieved 7 June 2022. National Review. December 2011. Archived from the original on 7 June 2022. National Review. December 2011. Archived from the original on 7 June 2022. National Review. December 2011. Archived from the original on 7 June 2022. National Review. December 2011. Archived from the original on 7 June 2022. National Review. December 2011. Archived from the original on 7 June 2022. National Review. December 2011. Archived from the original on 7 June 2022. National Review. December 2011. Archived from the original on 8 June 2022. National Review. December 2011. Archived from the original on 8 June 2022. National Review. December 2011. Archived from the original on 9 June 2022. National Review. December 2011. Archived from the original on 9 June 2022. National Review. December 2011. Archived from the original on 9 June 2022. National Review. December 2011. Archived from the original on 9 June 2022. National Review. December 2011. Archived from the original on 9 June 2022. National Review. December 2011. Archived from the original on 9 June 2022. National Review. December 2011. Archived from the original on 9 June 2022. National Review. December 2011. Archived from the original on 9 June 2022. National Review. December 2011. Archived from the original on 9 June 2022. National Review. December 2011. Archived from the original on 9 June 2022. National Review. December 2011. Archived from the original on 9 June 2022. National Review. December 2011. Archived from the original on 9 June 2022. National Review. December 2011. Archived from the original on 9 June 2022. National Review. December 2011. Archived from the original Review. December 2011. Archived from the original Review. December 2011. Archived from the orig
as a Marine Fuel Report (PDF) (Report). FC Business Intelligence. Archived (PDF) from the original on 10 April 2021. A Kamitani, A.; Morishita, S.; Kotaki, H.; Arscott, S. (2008)
 "Miniaturized microDMFC using silicon microsystems techniques: Performances at low fuel flow rates" (PDF). Journal of Micromechanics and Microengineering. 18 (12): 125019. Bibcode: 2008 MiMi..18l5019K. doi:10.1088/0960-1317/18/12/125019. S2CID 110214840. Archived (PDF) from the original on 21 November 2021. Retrieved 30 October
2021. ^ Kamitani, A.; Morishita, S.; Kotaki, H.; Arscott, S. (2011). "Microfabricated microfluidic fuel cells". Sensors and Actuators B: Chemical. 154 (2): 174. Bibcode: 2011SeAcB.154..174K. doi:10.1016/j.snb.2009.11.014. ^ Berger, Sandy (30 September 2006). "Methanol Laptop Fuel". Comput Kiss. Archived from the original on 3 February 2007.
Retrieved 22 May 2007. ^ Blum, Deborah (19 February 2010). "The little-told story of how the U.S. government poisoned alcohol during Prohibition". Slate Magazine. Archived from the original on 6 August 2018. Retrieved 10 June 2010. ^ Yant, W. P.; Schrenk, H. H.; Sayers, R. R. (1931). "Methanol Antifreeze and Methanol Poisoning". Industrial &
Engineering Chemistry. 23 (5): 551. doi:10.1021/ie50257a020. ^ Commission Regulation (EU) 2018/589 of 18 April 2018 amending Annex XVII to Regulation, Evaluation, Authorisation and Restriction of Chemicals (REACH) as regards methanol ^
Corrigendum to Commission Regulation (EU) 2018/589 of 18 April 2018 amending Annex XVII to Regulation (EU) No 1907/2006 of the European Parliament and of the Council concerning the Registration, Evaluation, Authorisation and Restriction of Chemicals (REACH) as regards methanol ^ "Annex XV Restriction Report: Proposal For a Restriction of Chemicals (REACH) as regards methanol ^ "Annex XV Restriction Report: Proposal For a Restriction
Substance Name: Methanol - table D.1-4-page 79". 16 January 2015. Archived from the original on 26 May 2021. Retrieved 29 November 2024. ^ Deutschmann, Olaf; Knözinger, Helmut; Kochloefl, Karl and Turek, Thomas (2012) "Heterogeneous Catalysis and Solid Catalysis, 3. Industrial Applications" in Ullmann's Encyclopedia of Industrial
Chemistry. Wiley-VCH, Weinheim. doi:10.1002/14356007.005 o03 Dozano, Giulia; Manenti, Flavio (2016). "Efficient methanol synthesis: Perspectives, technologies and optimization strategies". Progress in Energy and Combustion Science. 56: 76. Bibcode:2016PECS...56...71B. doi:10.1016/j.pecs.2016.06.001. ISSN 0360-1285. Mu-Hyun Baik;
Martin Newcomb; Richard A. Friesner; Stephen J. Lippard (2003). "Mechanistic Studies on the Hydroxylation of Methane by Methane Monooxygenase". Chem. Rev. 103 (6): 2385-2420. doi:10.1021/cr950244f. PMID 12797835. ^ a b Lawton, T. J.; Rosenzweig, A. C. (2016). "Biocatalysts for methane conversion: big progress on breaking a small
 substrate". Curr. Opin. Chem. Biol. 35: 142-149. doi:10.1016/j.cbpa.2016.10.001. PMC 5161620. PMID 27768948. ^ Alayon, E. M. C.; Nachtegaal, M.; Ranocchiari, M.; Ranocchiari, M.; Van Bokhoven, J. A. (2012). "Catalytic Conversion of Methane to Methanol Using Cu-Zeolites". CHIMIA International Journal for Chemistry. 66 (9): 668-674.
doi:10.2533/chimia.2012.668. PMID 23211724. Archived from the original on 26 May 2021. Retrieved 10 May 2021. A; Hutchings, R. L.; Dimitratos, N.; Lopez-Sanchez, J. A.; Ab Rahim, M. H.; Forde, M.M.; Thetford, A.; Murphy, D.M.; Hagen, H.; Stangland, E.E.; Moulijn, J.M.; Taylor, S. H.; Willock, D. J.; Hutchings, G.J. (2012). "Catalytic Archived from the original on 26 May 2021. A Retrieved 10 May 2021. The original on 26 May 2021. A Retrieved 10 May 2021. A Retrieved 1
Active Sites in Zeolites and Their Correlation to Metalloenzymes". Chemical Reviews. 118 (5): 2718-2768. doi:10.1021/acs.chemrev.7b00344. PMID 29256242. Archived from the original on 26 May 2021. Retrieved 25 September 2021. ^ Snyder, Benjamin E. R.; Vanelderen, Pieter; Bols, Max L.; Hallaert, Simon D.; Böttger, Lars H.; Ungur, Liviu;
 Pierloot, Kristine; Schoonheydt, Robert A.; Sels, Bert F. (2016). "The active site of low-temperature methane hydroxylation in iron-containing zeolites". Nature. 536 (7616): 317-321. Bibcode: 2016Natur.536..317S. doi:10.1038/nature19059. PMID 27535535. S2CID 4467834. ^ Stavrakou, T.; Guenther, A.; Razavi, A.; Clarisse, L.; Clerbaux, C.; Coheur.
P.-F.; Hurtmans, D.; Karagulian, F.; De Mazière, M.; Vigouroux, C.; Amelynck, C.; Schoon, N.; Laffineur, Q.; Heinesch, B.; Aubinet, M. (25 May 2011). "First space-based derivation of the global atmospheric methanol emission fluxes". Atmospheric Chemistry and Physics. 11 (10): 4873–4898. Bibcode: 2011ACP....11.4873S. doi:10.5194/acp-11-4873-4898.
2011. ISSN 1680-7324. S2CID 54685577. Archived from the original on 26 September 2022. Retrieved 26 September 2022. ^ a b c d e Sollai, Stefano; Porcu, Andrea; Tola, Vittorio; Ferrara, Francesca; Pettinau, Alberto (1 February 2023). "Renewable methanol production from green hydrogen and captured CO2: A techno-economic assessment".
 Journal of CO2 Utilization. 68: 102345. doi:10.1016/j.jcou.2022.102345. ISSN 2212-9820. Sasidhar, Nallapaneni (November 2023). "Carbon Neutral Fuels and Chemicals from Standalone Biomass Refineries" (PDF). Indian Journal of Environment Engineering. 3 (2): 1-8. doi:10.54105/jjee.81845.113223. ISSN 2582-9289. S2CID 265385618.
 Retrieved 3 December 2023. ^ "Green Hydrogen for Industry: A Guide to Policy Making". www.irena.org. 8 March 2022. p. 18. Retrieved 4 September 2023. ^ a b Vale A (2007). "Methanol Poisoning Overview". Antizol. Archived from the original on 5 October 2011. ^
 "Integrated Risk Information System". US EPA, ORD, NCEA, IRISD. 15 March 2013. Archived from the original on 5 December 2012. Retrieved 18 February 2013. ^ "Toxicological Review of Methanol (Noncancer) (CAS No. 67-56-1) In Support of Summary Information on the Integrated Risk Information System (IRIS)" (PDF). EPA. September 2013.
 EPA/635/R-11/001Fa. Archived (PDF) from the original on 31 March 2021. Retrieved 30 March 2021. Netrieved 30 March 2021. Schep LJ, Slaughter RJ, Vale JA, Beasley DM (2009). "A seaman with blindness and confusion". BMJ. 339: b3929. doi:10.1136/bmj.b3929. PMID 19793790
S2CID 6367081. Archived from the original on 8 October 2009. Retrieved 3 October 2009. ^ McMartin KE, Martin-Amat G, Noker PE, Tephly TR (1979). "Lack of a role for formaldehyde in methanol poisoning in the monkey". Biochem. Pharmacol. 28 (5): 645-9. doi:10.1016/0006-2952(79)90149-7. PMID 109089. ^ Liesivuori J, Savolainen H
(September 1991). "Methanol and formic acid toxicity: biochemical mechanisms". Pharmacol. Toxicol. 69 (3): 157-63, doi:10.1111/j.1600-0773.1991.tb01290.x. PMID 1665561. Peterson, Charles D. (1981). "Oral ethanol doses in patients with methanol poisoning". American Journal of Health-System Pharmacy. 38 (7): 1024-1027.
doi:10.1093/ajhp/38.7.1024. ^ Beauchamp, GA; Valento, M (September 2016). "Toxic Alcohol Ingestion: Prompt Recognition And Management in the Emergency Medicine Practice. 18 (9): 1-20. PMID 27538060. ^ Ferri, Fred F. (2016). Ferri's Clinical Advisor 2017: 5 Books in 1. Elsevier Health Sciences. p. 794. ISBN 978-0-
323-44838-3. Archived from the original on 10 January 2023. Retrieved 22 August 2019. ^ Kruse, JA (October 2012). "Methanol and ethylene glycol intoxication". Critical Care Clinics. 28 (4): 661-711. doi:10.1016/j.ccc.2012.07.002. PMID 22998995. ^ Clary, John J. (2013). The Toxicology of Methanol. John Wiley & Sons. p. 3.4.1. ISBN 978-1-118-
35310-3. Robert Boyle, The Sceptical Chymist (1661) - contains account of distillation of wood alcohol. Schmidt, Eckart W. (2022). "Methanol". Alcohols. Encyclopedia of Liquid Fuels. De Gruyter. pp. 2-12. doi:10.1515/9783110750287-001. ISBN 978-3-11-075028-7. International Chemical Safety Card 0057 Methyl Alcohol (Methanol) CDC/NIOSH,
links to safety information CDC - NIOSH Pocket Guide to Chemical Hazards - Methyl Alcohol Methanol Fact Sheet - National Pollutant Inventory Retrieved from 2 1-Pentanol Names Preferred IUPAC name Pentan-1-ol[1] Identifiers CAS Number 71-41-0 Y 3D model (JSmol) Interactive
image Beilstein Reference 1730975 ChEBI CHEBI:44884 Y ChEMBL 14568 Y ChemSpider 6040 Y ECHA InfoCard 100.000.684 EC Number 200-752-1 Gmelin Reference 25922 KEGG C16834 N MeSH n-Pentanol PubChem CID 6276 RTECS number SB9800000 UNII M9L931X26Y Y UN number 1105 CompTox Dashboard (EPA)
DTXSID6021741 InChI Inch
\log P = 1.348 \text{ Vapor pressure } 200 \text{ Pa (at } 20 \text{ °C)} \text{ Magnetic susceptibility } (\chi) = 67.7 \cdot 10 - 6 \text{ cm} 3/\text{mol} = 1.348 \text{ Vapor pressure } 200 \text{ Pa (at } 20 \text{ °C)} \text{ Magnetic susceptibility } (\chi) = 67.7 \cdot 10 - 6 \text{ cm} 3/\text{mol} = 1.348 \text{ Vapor pressure } 200 \text{ Pa (at } 20 \text{ °C)} \text{ Magnetic susceptibility } (\chi) = 67.7 \cdot 10 - 6 \text{ cm} 3/\text{mol} = 1.348 \text{ Vapor pressure } 200 \text{ Pa (at } 20 \text{ °C)} \text{ Magnetic susceptibility } (\chi) = 67.7 \cdot 10 - 6 \text{ cm} 3/\text{mol} = 1.348 \text{ Vapor pressure } 200 \text{ Pa (at } 20 \text{ °C)} \text{ Magnetic susceptibility } (\chi) = 67.7 \cdot 10 - 6 \text{ cm} 3/\text{mol} = 1.348 \text{ Vapor pressure } 200 \text{ Pa (at } 20 \text{ °C)} \text{ Magnetic susceptibility } (\chi) = 67.7 \cdot 10 - 6 \text{ cm} 3/\text{mol} = 1.348 \text{ Vapor pressure } 200 \text{ Pa (at } 20 \text{ °C)} \text{ Magnetic susceptibility } (\chi) = 67.7 \cdot 10 - 6 \text{ cm} 3/\text{mol} = 1.348 \text{ Vapor pressure } 200 \text{ Pa (at } 20 \text{ °C)} \text{ Magnetic susceptibility } (\chi) = 67.7 \cdot 10 - 6 \text{ cm} 3/\text{mol} = 1.348 \text{ Vapor pressure } 200 \text{ Pa (at } 20 \text{ °C)} \text{ Magnetic susceptibility } (\chi) = 67.7 \cdot 10 - 6 \text{ cm} 3/\text{mol} = 1.348 \text{ Vapor pressure } 200 \text{ Pa (at } 20 \text{ °C)} \text{ Magnetic susceptibility } (\chi) = 67.7 \cdot 10 - 6 \text{ cm} 3/\text{mol} = 1.348 \text{ Vapor pressure } 200 \text{ Pa (at } 20 \text{ °C)} \text{ Magnetic susceptibility } (\chi) = 67.7 \cdot 10 - 6 \text{ cm} 3/\text{mol} = 1.348 \text{ Vapor pressure } 200 \text{ Pa (at } 20 \text{ °C)} \text{ Magnetic susceptibility } (\chi) = 67.7 \cdot 10 - 6 \text{ cm} 3/\text{mol} = 1.348 \text{ Magnetic susceptibility } (\chi) = 67.7 \cdot 10 - 6 \text{ cm} 3/\text{mol} = 1.348 \text{ Magnetic susceptibility } (\chi) = 67.7 \cdot 10 - 6 \text{ cm} 3/\text{mol} = 1.348 \text{ Magnetic susceptibility } (\chi) = 67.7 \cdot 10 - 6 \text{ cm} 3/\text{mol} = 1.348 \text{ Magnetic susceptibility } (\chi) = 67.7 \cdot 10 - 6 \text{ cm} 3/\text{mol} = 1.348 \text{ Magnetic susceptibility } (\chi) = 67.7 \cdot 10 - 6 \text{ cm} 3/\text{mol} = 1.348 \text{ Magnetic susceptibility } (\chi) = 67.7 \cdot 10 - 6 \text{ cm} 3/\text{mol} = 1.348 \text{ Magnetic susceptibility } (\chi) = 67.7 \cdot 10 - 6 \text{ cm} 3/\text{mol} = 1.348 \text{ Magnetic susceptibility } (\chi) = 67.7 \cdot 10 - 6 \text{ cm} 3/\text{mol} = 1.348 \text{ Magne
mol-1 Hazards GHS labelling: Pictograms Signal word Warning Hazard statements H226, H315, H332, H335 Precautionary statements P261 NFPA 704 (fire diamond) 1 2 0 Flash point 49 °C (120 °F; 322 K) Autoignition temperature 300 °C (572 °F; 573 K) Related compounds Related compounds Hexane Pentylamine Except where otherwise noted,
data are given for materials in their standard state (at 25 °C [77 °F], 100 kPa). N verify (what is YN?) Infobox references Chemical compound 1-Pentanol, (or n-pentanol, pentan-1-ol), is an organic compound with the formula CH3CH2CH2CH2CH2OH and is classified as a primary alcohol. [2] It is a colourless liquid with a distinctive aroma. It is one of
8 isomeric alcohols with the formula C5H11OH. It is used as a solvent, a biological drying agent and in the synthesis of some fragrance compounds. It is also a common component of fusel alcohols (fusel oils), the undesirable byproducts of alcoholic fermentation. 1-Pentanol is prepared from 1-butene by hydroformylation followed by hydrogenation of
fermentation.[4][5] The hydroxyl group (OH) is the active site of many reactions. The ester formed from 1-pentanol and acetic acid is amyl acetate (also called pentyl acetate), which has a banana-like odor. It is a precursor to dipentyl zinc
dithiophosphates, which are used in froth flotation.[3] In 2014, a study was conducted comparing the performance of diesel fuel blends with various proportions of pentanol, particulate emissions decreased.[6] Pentanol is often used as a solvent. ^ "n-pentanol -
Compound Summary". PubChem Compound. USA: National Center for Biotechnology Information. 26 March 2005. Identification and Related Records, Retrieved 10 October 2011. ^ CRC Handbook of Chemistry and Physics 65th ed. ^ a b Lappe, Peter; Hofmann, Thomas (2011). "Pentanols". Ullmann's Encyclopedia of Industrial Chemistry.
doi:10.1002/14356007.a19 049.pub2. ISBN 9783527303854. ^ Cann, Anthony F.; Liao, James C. (2010-01-01). "Pentanol isomer synthesis in engineered microorganisms". Applied Microbiology and Biotechnology. 85 (4): 893-899. doi:10.1007/s00253-009-2262-7. ISSN 1432-0614. PMC 2804790. PMID 19859707. ^ Tseng, Hsien-Chung (2011).
Production of pentanol in metabolically engineered Escherichia coli (Thesis thesis). Massachusetts Institute of Technology. hdl:1721.1/65767. Wei, Liangjie & Cheung, C.s & Huang, Zuohua. (2014). Effect of n-pentanol addition on the combustion, performance and emission characteristics of a direct-injection diesel engine. Energy. 70.
10.1016/j.energy, 2014.03.106. Retrieved from 3 The following pages link to 1-Pentanol External tools (links | edit) Benzodiazepine (links | edit) Benzodiazepine (links | edit) Amyl alcohol (chemistry) (links | edit) Amyl alcohol (links | edit) Benzodiazepine (links | edit)
Cholesterol (links | edit) Carbamazepine (links | edit) Ethanol (links | edit) Flunitrazepam (links | edit) Nicotinamide (links | edit) Nicoti
Butanol (links | edit) Sorbitol (links | edit) Progesterone (links | edit) Acetophenone (links | edit) Mochi (links | edit) Moch
oxide (links | edit) Temazepam (links | edit) Alprazolam (links | edit) Lorazepam (links | edit) Lorazepam (links | edit) Dichloromethane (links | edit) Di
edit) Menthol (links | edit) Isoflurane (lin
```